

Prevention of catheter-associated UTI

Paul Anantharajah Tambyah¹

Leyland Chuang²

Edmund Chiong³

¹Department of Medicine, Yong Loo Lin School of Medicine, National University Hospital, Singapore, Singapur

²Department of Medicine, Ng Teng Fong Hospital, National University Health System, Singapore, Singapur

³Department of Urology, National University Hospital, Singapore

Abstract

We surveyed existing guidelines and literature on the prevention of catheter-associated urinary tract infection (CAUTI). Key principles include avoiding unnecessary urinary catheterization, maintaining a closed aseptic drainage system and minimizing duration of catheterization. Institutional guidelines on appropriate indications for catheterization, reminder and stop-order systems, and policies on restricted urinary catheterization and nurse-directed catheter removal are effective and recommended. Urinary catheter care bundles and infection control interventions described in various guidelines are summarized and should be considered. While routine use of anti-infective coated catheters are not recommended, anti-infective-impregnated catheters may reduce risks of CAUTI for short term catheterization, and may be considered if shown to be cost effective. Evidence for the use of routine systemic antibiotic prophylaxis for the prevention of CAUTI is inconsistent and insufficient. Treatment of symptomatic CAUTI should be guided by local uropathogens, susceptibility patterns and urine culture results, and antibiotic should be deescalated to the narrowest spectrum choice where possible. Routine treatment of catheter-associated asymptomatic bacteriuria is not recommended.

Summary of recommendations

Recommendation	LE/GR
Avoiding unnecessary catheterization and prompt removal	
1. Establishing institutional guidelines to prevent inappropriate urinary catheterization has been shown to reduce catheter use and CAUTI rates [1], [2], [3], [4].	2b/B
2. Early removal of IUCs reduces the risks of subsequent CAUTI [1], [3], [5], [6] and other complications [5], [7].	2b/B
3. Health care workers are often unaware about the presence of IUCs in patients [8], [9]. This should be addressed, in order to minimize inappropriate use of IUCs and reduce duration of catheterization.	3/B
4. Nurse generated or electronic-based reminder and stop-order systems for the removal of urinary catheters reduce utilization of IUCs and rates of CAUTIs [1], [3], [4], [10], [11], [12], [13], [14], [15], [16], [17], [18], [19], [20], [21], [22], [23], [24].	1a/B
5. A restricted urinary catheterization policy with daily chart review of appropriateness of catheter insertion reduces urinary catheterization rates [25].	2b/B
6. Nurse-directed urinary catheter removal reduces urinary catheter utilization and rates of CAUTI [16], [20], [22], [26], [27].	2a/B

Recommendation **LE/GR**

Avoiding unnecessary catheterization and prompt removal

- 7. Alternative methods of bladder drainage such as SPCs and intermittent catheterization may be considered, but evidence of efficacy in preventing symptomatic UTI remains limited [28], [29], [30], [31]. 1b/C
- 8. Educational interventions may reduce IUC utilization [2], [4], [32]. 2a/B

Care and maintenance of urinary catheter system

- 9. Maintaining sterility of closed urinary catheter systems is recommended to prevent the development of CAUTIs [33], [34], [35], [36], [37], [38], [39], [40]. 3/B

Urinary catheter types

Evidence for various urinary catheter types are as follows:

- No recommendations on the use of hydrophilic-coated catheters for intermittent catheterization can be made [41], [42], [43]. 1a
- 10.
 - Anti-infective-impregnated catheters reduce risks of CAUTI slightly in short-term catheterization of 14 days or less, and may be cost effective [44], [45], [46]. 1b/B
 - Silver alloy catheters reduce risks of bacteriuria, but not CAUTI, and are unlikely to be cost effective [44], [45], [46]. 1a/A
 - There are insufficient data to recommend the use of antibiotic-impregnated or silver alloy catheters for long-term catheterization [47]. 1b

CAUTI prophylaxis

No recommendation can be made for the routine use of antibiotic prophylaxis for the prevention of CAUTI due to inconsistent and insufficient evidence, and potential risks of increased antimicrobial resistance. It may be considered where benefits are deemed to outweigh risks.

- 11.
 - Use of antibiotic prophylaxis during short-term urinary catheterization of up to 14 days in adult patients is associated with significant reduction in asymptomatic bacteriuria, but there is no strong evidence showing reduction in symptomatic UTIs [48], [49]. 1a
 - Antibiotic prophylaxis at time of short-term IUC removal is associated with significant reduction in symptomatic UTI, but cannot be recommended routinely due to high numbers needed to treat, and the potential to increase antimicrobial resistance rates [50], [51].
 - There is insufficient evidence to recommend antimicrobial prophylaxis for long-term catheterization of more than 14 days [30].

Treatment of CAUTI

- 12. Empiric antimicrobial therapy may be guided by recent prior urine culture results, where possible [52]. Catheter removal and if necessary reinsertion remains the key to treatment. 2b/C

Recommendation	LE/GR
Avoiding unnecessary catheterization and prompt removal	
13. Early de-escalation of antibiotic therapy, as guided by urine culture results, to the narrowest spectrum antibiotic available.	4/C
14. Shorter 5 day course of antibiotics with catheter exchange may be considered in the treatment of CAUTI in patients with spinal cord injury [53].	1b/B
15. Catheter-associated asymptomatic bacteriuria should not be routinely treated with antibiotics [35], [54].	1b/A

1 Introduction

Urinary tract infections (UTIs) are the commonest cause of nosocomial infections worldwide, and have been estimated to cause approximately 30% of healthcare-associated infections in the acute care setting in the United States [55]. Approximately 75% of healthcare-associated UTIs are associated with an indwelling urinary catheter (IUC), and 12 to 16 percent of hospitalized patients will undergo urinary catheterization [56]. The high frequency of IUC utilization in healthcare highlights the impact and challenge of catheter-associated UTI (CAUTI) in the healthcare system globally [57], [58], [59], [60], [61], [62], [63] (LE: 3).

The aim of this chapter is to highlight key evidence-based recommendations for the prevention of CAUTIs.

2 Methods

Current published major practice guidelines were reviewed, and their recommendations were summarized. A systematic literature search of studies from the past ten years was also performed in PubMed with the following key words: urinary tract infection, bacteriuria, CAUTI, UTI, catheter. Search criteria were limited by following: availability of abstracts, English language and adults. Publications identified were screened by title and abstracts for relevance, and supplemented by citations and publications known to the authors. A total of 120 publications were included into the review.

The studies were rated according to the level of evidence and the strength of recommendations graded according to the system used in the EAU guidelines (2015) modified from the Oxford Centre for Evidence-based Medicine [37].

3 Pathogenesis and risk factors

The presence of a transurethral catheter predisposes patients to CAUTIs by bypassing or inhibiting natural host defenses [64]. The development of biofilm on the urinary catheters further exacerbates this by providing a favorable environment for bacterial proliferation and invasion [38], [65] (LE: 3).

Bacteria may be introduced into the urinary tract via multiple routes:

- i. Inoculation during catheter insertion (e.g. non-compliance to aseptic technique during catheter insertion).
- ii. Intraluminal ascent after contamination of the closed urinary catheter system. This may be related to lapses in aseptic practices while emptying the urinary drainage bag, or after disconnection of catheters from urinary bags.
- iii. Extraluminal route of ascent along the external catheter surface into the urethra.

Risk factors for CAUTIs which have been identified in prospective observational studies include [64], [66], [67], [68] (LE: 2b):

- Duration of catheterization
- Poor catheter care or breaks in aseptic technique
- Female gender
- Diabetes mellitus

- Anatomical or functional abnormalities of the urinary tract
- Insertion of the catheter outside the operating room

In particular, longer duration of catheterization was highlighted as a risk factor for CAUTI by Barbadoro et al., who found catheterization of greater than 4 days to be associated with CAUTI (odds ratio 8.21; 95% confidence interval, 3.79–17.73; $p < 0.05$) [68].

4 Summary of current guidelines on CAUTI prevention

Various guidelines on prevention of CAUTI have been published. Current guidelines include those from the Healthcare Infection Control Practices Advisory Committee (HICPAC) [33], the Infectious Diseases Society of America (IDSA) [34], the Society for Healthcare Epidemiology of America (SHEA) [35], the Department of Health of England (epic3 guidelines) [36], the European Association of Urology (EAU) [37], and also the joint European and Asian guidelines [38]. Their recommendations are summarized in table 1.

Table 1: Summary of recommendations from published guidelines

	European & Asian guidelines 2008 [38]	IDSA 2009 [34]	CDC/HICPAC 2009 [33]	epic3 2014 [36]	SHEA/IDSA 2014 [35]	EAU 2015 [37]
General						
Provide guidelines on catheter use, insertion and maintenance	Y	Y	Y	Y	Y	Y
Documentation of urinary catheterization: may include details of catheter insertion and removal, indication for insertion, and daily maintenance care tasks	ND	Y	Y	Y	Y	ND
Educate and train healthcare personnel	ND	Y	Y	Y	Y	ND
Hand hygiene compliance during patient care	Y	ND	Y	Y	Y	Y
Catheter insertion						
Evaluate need for catheterization						
Consider necessity of catheterization, ensure catheterization only for appropriate indications	ND	Y	Y	Y	Y	ND
Consider alternatives (e.g. condom catheters, intermittent straight catheterization and suprapubic catheters)	Y	Y	Y	Y	Y	Y
Protocol for post-operative urinary retention (e.g. intermittent catheterization, use of bladder scanners)	ND	Y	Y	ND	Y	ND
Choice of catheter						
Use smallest gauge urinary catheter possible	Y	ND	Y	Y	Y	Y
Choice of standard catheter material type	U	ND	Y	U	ND	ND
Routine use of antimicrobial/antiseptic-impregnated catheters is not recommended	Y	Y	U	ND	Y	Y
Routine use of silver alloy catheters is not recommended	Y	Y	U	ND	ND	U
Use of hydrophilic catheters for intermittent catheterization recommended	ND	ND	Y	ND	ND	ND

	European & Asian guidelines 2008 [38]	IDSA 2009 [34]	CDC/HICPAC 2009 [33]	epic3 2014 [36]	SHEA/IDSA 2014 [35]	EAU 2015 [37]
Use of pre-connected urinary system	ND	Y	Y	ND	ND	ND
Catheter insertion						
Use aseptic technique and sterile equipment	Y	Y	Y	Y	Y	Y
Use adequate lubricant	Y	ND	Y	Y	Y	Y
Antiseptic solution for cleaning of meatus during catheter insertion	U	ND	U	N	Y	ND
Barrier precaution	ND	U	Y	ND	Y	ND
Insertion of catheter by trained personnel	ND	Y	Y	Y	Y	ND
Train patients and family	ND	ND	Y	Y	ND	ND
Catheter maintenance						
Catheter review						
Minimize duration of catheterization	Y	Y	Y	Y	Y	Y
Regular review of ongoing need for catheter	ND	ND	Y	Y	Y	ND
System to identify and remove catheters that are no longer necessary: may include electronic or nursing reminders, automatic stop orders, or nurse led catheter removal protocols	ND	Y	Y	Y	Y	Y
Early removal of IUCs post-operatively	Y	ND	Y	ND	ND	Y
Catheter care						
Maintain closed drainage system	Y	Y	Y	Y	Y	Y
Replace system if there is any break in asepsis	ND	ND	Y	ND	Y	ND
No routine change in catheter	Y	U	Y	Y	Y	Y
Obtain urine samples aseptically	ND	ND	Y	Y	Y	ND
Routine hygiene for meatal care. Topical antiseptic/antibiotic applied to catheter, urethra or meatus are not recommended.		Y	Y	Y	Y	Y
Secure catheter	ND	ND	Y	Y	Y	ND
Avoid irrigation for purpose of preventing infection	ND	Y	Y	Y	Y	Y
Maintain unobstructed urine flow	ND	ND	Y	ND	Y	ND
Keep collecting bag below level of bladder	ND	Y	Y	Y	Y	Y
Keep collecting bag off the floor	ND	ND	Y	Y	Y	ND
Empty collecting bag regularly using separate collecting container for each patient, avoid touching the draining spigot to the container	ND	ND	Y	Y	Y	ND
Separate patients with catheters	ND	U	U	ND	U	ND
Asymptomatic bacteriuria						

	European & Asian guidelines 2008 [38]	IDSA 2009 [34]	CDC/HICPAC 2009 [33]	epic3 2014 [36]	SHEA/IDSA 2014 [35]	EAU 2015 [37]
Do not routinely screen for bacteriuria in asymptomatic catheterized patients	Y	Y	Y	ND	Y	Y
Do not routinely treat asymptomatic bacteriuria in catheterized patients	Y	Y	ND	ND	Y	Y
Antiseptic & antimicrobial prophylaxis						
Routine use of systemic antimicrobial prophylaxis not recommended						
Short-term catheterization	ND	Y	Y	ND	Y	Y
Intermittent catheterization	Y					Y
Long-term catheterization	U					U
Routine use of long term antibiotic suppressive therapy not recommended	Y	ND	ND	ND	ND	Y
Routine use of urinary antiseptics (e.g. methenamine) not recommended	Y	Y	U	ND	U	Y
Routine addition of antiseptic or antimicrobial solutions to urinary drainage bags not recommended	ND	Y	Y	Y	ND	ND
Surveillance & audit						
Performance feedback to clinical staff	ND	Y	Y	Y	Y	ND
Surveillance: may include CAUTI rates and catheter utilization rates	ND	ND	Y	ND	Y	ND

Legend: Y: Recommended; N: Not recommended; U: Unresolved; ND: Not discussed

5 Recommendations for CAUTI prevention

5.1 Principles of CAUTI prevention

Principles for prevention of CAUTI may be broadly classified under the following categories:

- Avoiding unnecessary urinary catheterization and minimizing duration of catheterization;
- Preserving closed aseptic drainage of the urinary catheter system;
- Implementing urinary catheter care bundles and infection prevention programs.

In an American multicenter prospective surveillance study, Lewis et. al. found that 72% of CAUTIs occurred in patients not in intensive care units [69]. Hence, CAUTI prevention efforts should include patients in general wards.

5.2 Avoiding unnecessary catheterization and prompt removal

Establishing institutional guidelines to prevent inappropriate urinary catheterization reduces catheter use and CAUTI rates [1], [2], [3], [4] (LE/GR: 2b/B). Indications for catheterization which are considered appropriate by expert panels include the following [33], [35], [36], [70] (LE/GR: 4/B):

1. Acute urinary retention or bladder outlet obstruction;
2. Management of open sacral or perineal wounds in incontinent patients;
3. Urinary incontinence in patients for whom nurses find it difficult to provide skin care despite other urinary management strategies (e.g. hemodynamic or respiratory instability with movement, conditions requiring strict prolonged immobility);
4. Frequent measurement (e.g. in critically ill patients) and/or precise measurement of urine volumes required to provide treatment, that cannot be assessed by other strategies;
5. Single 24 hour urine sample collection for diagnostic tests that cannot be obtained by other strategies;
6. Reduce acute, severe pain with movement when other urine management strategies are difficult;
7. Improvement in comfort in end of life care if needed;
8. Management of gross hematuria with blood clots in urine;
9. Clinical conditions where non-indwelling strategies are inadequate, or where intermittent straight catheterization or external catheter placement are difficult;
10. Perioperative use in selected surgical procedures (e.g. urologic surgery, prolonged duration of surgery).

Reduction in utilization of IUCs and their early removal have been shown to reduce the risks of CAUTI [[1](#) [3](#), [5](#), [6](#)] and other complications, such as mortality [[5](#)] and bladder cancers [[7](#)] (LE/GR: 2b/B). A Cochrane study on early removal in patients with short-term IUCs showed lower risks of UTI and shorter hospitalization [[71](#)]. However, healthcare workers are often unaware about the presence of IUCs in patients [[8](#), [9](#)], contributing to unnecessary or inappropriate catheterization (LE/GR: 3/B). This may be prevented by nurse generated or electronic-based reminders and stop-order systems for the removal of urinary catheters, which have been shown in multiple studies and a systematic review to reduce utilization of IUCs and rates of CAUTIs [[1](#), [3](#), [4](#), [10](#), [11](#), [12](#), [13](#), [14](#), [15](#), [16](#), [17](#), [18](#), [19](#), [20](#), [21](#), [22](#), [23](#), [24](#)] (LE/GR: 1a/B). A restricted urinary catheterization policy together with daily chart review of appropriateness of new catheter insertions has also been shown to reduce catheterization rates [[25](#)] (LE/GR: 2b/B). Nurse-directed urinary catheter removal has also been shown to reduce catheter utilization and CAUTI rates [[16](#), [20](#), [22](#), [26](#), [27](#)] (LE/GR: 2a/B).

Alternative methods of bladder drainage other than IUCs may be considered, but evidence of efficacy in preventing CAUTI is limited (LE/GR: 1b/C). Recent trials were directed mostly at the use of suprapubic catheters (SPCs) and intermittent catheterization. A Cochrane review by Kidd et al. on short-term catheterization (of ≤14 days) in hospitalized adults found no conclusive evidence of reduction in symptomatic UTI with SPC compared to IUCs (LE: 1a), while results comparing IUC against intermittent catheterization were inconclusive and of low quality [[28](#)]. Two Cochrane reviews, one on alternative methods of long-term management of neurogenic bladder, and another comparing IUCs against intermittent and suprapubic catheterization for long term bladder drainage, could find no definitive evidence from randomized trials [[29](#), [30](#)]. A review by Hunter et al. found five non-randomized or retrospective studies showing no evidence of difference in UTI prevalence between IUC and SPC management of long term bladder drainage [[31](#)]. Despite these, there may be some evidence of benefit in reduction of asymptomatic bacteriuria and pain with the use of SPC [[28](#)] (LE: 1a), and more rapid return to normal micturition after urogynaecological surgery (LE: 1b) and acute urinary retention (LE: 2a) with the use of intermittent catheterization compared to IUCs [[72](#), [73](#)]. The weighing of incontinence pads for monitoring of urine output has also been shown to allow the removal of IUCs used for those purposes [[74](#)].

Education interventions (alone or as part of a bundle) are associated with reduction in IUC utilization, and may reduce inappropriate antimicrobial therapy for asymptomatic bacteriuria and CAUTI rates [[2](#), [4](#), [32](#)] (LE/GR: 2a/B).

Recommendation	LE/GR
1. Establishing institutional guidelines to prevent inappropriate urinary catheterization has been shown to reduce catheter use and CAUTI rates [1 , 2 , 3 , 4].	2b/B
2. Early removal of IUCs reduces the risks of subsequent CAUTI [1 , 3 , 5 , 6] and other complications [5 , 7].	2b/B
3. Health care workers are often unaware about the presence of IUCs in patients [8 , 9]. This should be addressed, in order to minimize inappropriate use of IUCs and reduce duration of catheterization.	3/B

4. Nurse generated or electronic-based reminder and stop-order systems for the removal of urinary catheters reduce utilization of IUCs and rates of CAUTIs [1], [3], [4], 1a/B [10], [11], [12], [13], [14], [15], [16], [17], [18], [19], [20], [21], [22], [23], [24].
5. A restricted urinary catheterization policy with daily chart review of appropriateness of catheter insertion reduces urinary catheterization rates [25]. 2b/B
6. Nurse-directed urinary catheter removal reduces urinary catheter utilization and rates of CAUTI [16], [20], [22], [26], [27]. 2a/B
7. Alternative methods of bladder drainage such as SPCs and intermittent catheterization may be considered, but evidence of efficacy in preventing symptomatic UTI remains limited [28], [29], [30], [31]. 1b/C
8. Educational interventions may reduce IUC utilization [2], [4], [32]. 2a/B

5.3 Care and maintenance of urinary catheter system

Maintaining sterility of urinary catheter systems is recommended, with all current guidelines reviewed above recommending aseptic catheterization technique and maintenance of closed drainage system [33], [34], [35], [36], [37], [38] (LE/GR: 3/B). Data from the Spanish national database of hospital infections and a Japanese multicenter prospective observational study showed closed urinary drainage systems were associated with significant reductions in CAUTI incidence [39], [40] (LE: 3). However, three randomized controlled trials examining the effect of pre-connected or sealed urinary catheter drainage systems on preventing bacteriuria showed differing outcomes [75], [76], [77] (LE: 1b). Also, a Cochrane systematic review [41] and multiple randomized controlled trials [78], [79], [80] comparing aseptic against clean techniques of catheterization have not shown differences in CAUTI rates (LE: 1a). All these studies however, had closed drainage in all systems.

The use of catheter securing devices has not been shown to reduce UTI rates [81] (LE: 1b), but should be considered to prevent catheter movement and urethral traction. Bladder irrigation [82], [83] (LE: 1b), antimicrobial additives to drainage bags [84] (LE: 1b), perineal cleansing with topical antiseptic agents [85], [86] (LE: 1b), daily bath with chlorhexidine washcloths [87], [88], [89] (LE: 2a) and complicated urinary drainage systems (e.g. anti-reflux valves and antiseptic release cartridge) [90], [91] (LE: 1b) have also not been shown to prevent CAUTI or bacteriuria and are not recommended (LE/GR: 1b/A). There is insufficient evidence to make recommendations on optimal time between catheter replacements [92].

Recommendation	LE/GR
1. Maintaining sterility of closed urinary catheter systems is recommended to prevent the development of CAUTIs [33], [34], [35], [36], [37], [38], [39], [40].	3/B

5.4 Urinary catheter types

Large numbers of recent studies have looked at the use of various catheter materials for the prevention of CAUTI, and anti-infective-impregnated catheters may be considered in some settings if cost-effective.

Hydrophilic-coated catheters have been studied mainly for intermittent catheterization in patients with spinal cord injury (SCI) or neurogenic bladder, and conclusions by three systematic reviews were inconsistent. The reviews for The Cochrane Collaboration [41] and National Institute for Health and Clinical Excellence (NICE) [42] did not show reduction in CAUTI, while Li et al. [43] found significantly lower incidence of UTIs. As such, no recommendation can be made (LE: 1a).

A Cochrane review on short-term catheterization (of ≤14 days) in adult hospitalized patients showed slight reduction in risks of CAUTI with the use of nitrofurazone-impregnated catheters, and slight reduction in risks of bacteriuria but not symptomatic CAUTI with silver alloy catheters [44] (LE: 1a). Trials with economic analysis funded by the United Kingdoms (UK) National Institute for Health Research found that the chances of cost-effectiveness are greater than 70% for nitrofurazone-impregnated catheters, and very unlikely for silver alloy catheters, in the setting of the UK National Health Service [45], [46]. This may not apply to all health systems.

The Cochrane review by Jahn et al. on IUC choice for long-term bladder drainage (>30 days) in adults was unable to find sufficient quality trials to provide reliable recommendations [47] (LE: 1b).

Trials exploring the use of various novel catheter coatings with other anti-septics [93], [94], [95], [96], [97] [98], [99], [100], non-pathogenic *Escherichia coli* [101], [102], [103], anti-biofilm agents [104] and anti-quorum sensing agents [105], [106] have been published, and await further confirmatory clinical outcome results.

Recommendation	LE/GR
Evidence for various urinary catheter types are as follows:	
1. <ul style="list-style-type: none"> • No recommendations on the use of hydrophilic-coated catheters for intermittent catheterization can be made [41], [42], [43]. • Anti-infective-impregnated catheters reduce risks of CAUTI slightly in short-term catheterization of 14 days or less, and may be cost effective [44], [45], [46]. • Silver alloy catheters reduce risks of bacteriuria, but not CAUTI, and are unlikely to be cost effective [44], [45], [46]. • There are insufficient data to recommend the use of antibiotic-impregnated or silver alloy catheters for long-term catheterization [47]. 	1a 1b/B 1a/A 1b

5.5 CAUTI prophylaxis

Multiple systematic reviews have examined the use of antibiotic prophylaxis in CAUTI prevention. Lusardi et al. examined adults on short-term catheters (≤ 14 days) [48], while Morton et al. examined persons with neurogenic bladder due to SCI [49], of whom many were on intermittent catheterization. Both studies found significant reduction in asymptomatic bacteriuria, but no strong evidence of reduction in symptomatic UTIs. In addition, Morton et al. found 4 of 5 trials reporting increase in proportion of antimicrobial resistance bacteria.

Marschall et al. performed a meta-analysis on the effects of antibiotic prophylaxis given upon removal of short-term IUC, and found a 5.8% absolute risk reduction in symptomatic UTI (number needed to treat to prevent one symptomatic UTI of 17) [50] (LE: 1a). However, the study authors and other reviews [51] have expressed reservations about routine antimicrobial prophylaxis upon catheter removal, in light of the likely increase in antibiotic consumption and its association with increased antimicrobial resistance. More research is needed to identify patient groups who will best benefit from this, such as kidney transplant recipients after transplant surgery [110] (LE: 2a).

Niël-Weise et al. performed a systematic review on antimicrobial prophylaxis for long-term catheterization of greater than 14 days, but data were sparse and outcomes inconsistent (LE: 1b) [30]. One observational prospective study examined prophylaxis with weekly high dose oral cycling antibiotic in adult patients with SCI and recurrent UTIs who are on long-term clean intermittent catheterization [111]. It found significant reduction in frequency of UTIs, mean duration of antimicrobial therapy for UTIs, and hospitalization days over a median follow-up duration of 29 months (LE: 2b), but routine use for long term prophylaxis remains controversial.

Studies on cranberry extract [112], [113] and oral vinegar [114] found no evidence of effect on CAUTI prevention (LE: 1b). Methenamine hippurate may have some benefit in UTI prevention in patients without renal tract abnormalities (LE: 1a) [115], and bacterial interference may reduce risk of UTI in patients with SCI (LE: 1b) [116], [117].

Recommendation	LE/GR
No recommendation can be made for the routine use of antibiotic prophylaxis for the prevention of CAUTI due to inconsistent and insufficient evidence, and potential risks of increased antimicrobial resistance. It may be considered where benefits are deemed to outweigh risks.	

1.
 - Use of antibiotic prophylaxis during short-term urinary catheterization of up to 14 days in adult patients is associated with significant reduction in asymptomatic bacteriuria, but there is no strong evidence showing reduction in symptomatic UTIs [48], [49]. 1a
 - Antibiotic prophylaxis at time of short-term IUC removal is associated with significant reduction in symptomatic UTI, but cannot be recommended routinely due to high numbers needed to treat, and the potential to increase antimicrobial resistance rates [50], [51].
 - There is insufficient evidence to recommend antimicrobial prophylaxis for long-term catheterization of more than 14 days [30].

6 Treatment of CAUTI

Choice of empiric antibiotic therapy may be guided by local uropathogens susceptibility patterns, and also recent prior urine culture results [52] (LE/GR: 2b/C). This should be de-escalated to culture guided narrow-spectrum antibiotics when possible (LE/GR: 4/C). Duration of treatment remains controversial, and guideline recommendations vary from 5 to 21 days [34], [38], [118], [119]. A shorter 5 day course of antibiotics with catheter exchange may be considered in the treatment of CAUTI in patients with SCI [53] (LE/GR: 1b/B). Routine treatment of catheter-associated asymptomatic bacteriuria is not recommended (LE/GR: 1b/B), with the exception of pregnancy and prior to traumatic genitourinary procedures [34], [54], [120].

Recommendation	LE/GR
1. Empiric antimicrobial therapy may be guided by recent prior urine culture results, where possible [52]. Catheter removal and if necessary reinsertion remains the key to treatment.	2b/C
2. Early de-escalation of antibiotic therapy, as guided by urine culture results, to the narrowest spectrum antibiotic available.	4/C
3. Shorter 5 day course of antibiotics with catheter exchange may be considered in the treatment of CAUTI in patients with spinal cord injury [53].	1b/B
4. Catheter-associated asymptomatic bacteriuria should not be routinely treated with antibiotics [35], [54].	1b/A

Conflicts of interest

Paul Anantharajah Tambyah has received research support from GlaxoSmithKline, Sanofi, Fab'entech, Teleflex, and Inviragen.

Edmund Chiong has had advisory roles too, and received conference support from Bayer, Janssen, GlaxoSmithKline, and Sanofi.

Leyland Chuang declares no conflict of interest.

Glossary

CAUTI: Catheter-associated urinary tract infection

EAU: European Association of Urology

HICPAC: Healthcare Infection Control Practices Advisory Committee (HICPAC)

IDSA: Infectious Diseases Society of America

IUC: Indwelling urinary catheter

NICE: National Institute for Health and Clinical Excellence

SCI: Spinal cord injury

SHEA: Society for Healthcare Epidemiology of America

SPC: Suprapubic catheters

UK: United Kingdoms

UTI: Urinary tract infection

References

1. Elpern EH, Killeen K, Ketchem A, Wiley A, Patel G, Lateef O. Reducing use of indwelling urinary catheters and associated urinary tract infections. *Am J Crit Care*. 2009 Nov;18(6):535-41; quiz 542. DOI: [10.4037/ajcc2009938](https://doi.org/10.4037/ajcc2009938)
2. Fakih MG, Pena ME, Shemes S, Rey J, Berriel-Cass D, Szpunar SM, Savoy-Moore RT, Saravolatz LD. Effect of establishing guidelines on appropriate urinary catheter placement. *Acad Emerg Med*. 2010 Mar;17(3):337-40. DOI: [10.1111/j.1553-2712.2009.00677.x](https://doi.org/10.1111/j.1553-2712.2009.00677.x)
3. Rothfeld AF, Stickley A. A program to limit urinary catheter use at an acute care hospital. *Am J Infect Control*. 2010 Sep;38(7):568-71. DOI: [10.1016/j.ajic.2009.12.017](https://doi.org/10.1016/j.ajic.2009.12.017)
4. Egger M, Balmer F, Friedli-Wüthrich H, Mühlemann K. Reduction of urinary catheter use and prescription of antibiotics for asymptomatic bacteriuria in hospitalised patients in internal medicine: before-and-after intervention study. *Swiss Med Wkly*. 2013 May 27;143:w13796. DOI: [10.4414/smw.2013.13796](https://doi.org/10.4414/smw.2013.13796)
5. Wald HL, Ma A, Bratzler DW, Kramer AM. Indwelling urinary catheter use in the postoperative period: analysis of the national surgical infection prevention project data. *Arch Surg*. 2008 Jun;143(6):551-7. DOI: [10.1001/archsurg.143.6.551](https://doi.org/10.1001/archsurg.143.6.551)
6. Al-Hazmi H. Role of duration of catheterization and length of hospital stay on the rate of catheter-related hospital-acquired urinary tract infections. *Res Rep Urol*. 2015;7:41-7. DOI: [10.2147/RRU.S75419](https://doi.org/10.2147/RRU.S75419)
7. West DA, Cummings JM, Longo WE, Virgo KS, Johnson FE, Parra RO. Role of chronic catheterization in the development of bladder cancer in patients with spinal cord injury. *Urology*. 1999 Feb;53(2):292-7. DOI: [10.1016/S0090-4295\(98\)00517-2](https://doi.org/10.1016/S0090-4295(98)00517-2)
8. Saint S, Wiese J, Amory JK, Bernstein ML, Patel UD, Zemencuk JK, Bernstein SJ, Lipsky BA, Hofer TP. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med*. 2000 Oct 15;109(6):476-80. DOI: [10.1016/S0002-9343\(00\)00531-3](https://doi.org/10.1016/S0002-9343(00)00531-3)
9. Harley G, Yeo AL, Stuart RL, Dendle C. A real-life snapshot of the use and abuse of urinary catheters on general medical wards. *Infect Control Hosp Epidemiol*. 2011 Dec;32(12):1216-8. DOI: [10.1086/662625](https://doi.org/10.1086/662625)
10. Meddings J, Rogers MA, Macy M, Saint S. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis*. 2010 Sep;51(5):550-60. DOI: [10.1086/655133](https://doi.org/10.1086/655133)
11. Apisarnthanarak A, Thongphubeth K, Sirinvaravong S, Kitkangvan D, Yuekyen C, Warachan B, Warren DK, Fraser VJ. Effectiveness of multifaceted hospitalwide quality improvement programs featuring an intervention to remove unnecessary urinary catheters at a tertiary care center in Thailand. *Infect Control Hosp Epidemiol*. 2007 Jul;28(7):791-8. DOI: [10.1086/518453](https://doi.org/10.1086/518453)
12. Crouzet J, Bertrand X, Venier AG, Badoz M, Husson C, Talon D. Control of the duration of urinary catheterization: impact on catheter-associated urinary tract infection. *J Hosp Infect*. 2007 Nov;67(3):253-7. DOI: [10.1016/j.jhin.2007.08.014](https://doi.org/10.1016/j.jhin.2007.08.014)
13. Robinson S, Allen L, Barnes MR, Berry TA, Foster TA, Friedrich LA, Holmes JM, Mercer S, Plunkett D, Vollmer CM, Weitzel T. Development of an evidence-based protocol for reduction of indwelling urinary catheter usage. *Medsurg Nurs*. 2007 Jun;16(3):157-61.
14. Huang WC, Wann SR, Lin SL, Kunin CM, Kung MH, Lin CH, Hsu CW, Liu CP, Lee SS, Liu YC, Lai KH, Lin TW. Catheter-associated urinary tract infections in intensive care units can be reduced by prompting physicians to remove unnecessary catheters. *Infect Control Hosp Epidemiol*. 2004 Nov;25(11):974-8. DOI: [10.1086/502329](https://doi.org/10.1086/502329)
15. Bruminhent J, Keegan M, Lakhani A, Roberts IM, Passalacqua J. Effectiveness of a simple

- intervention for prevention of catheter-associated urinary tract infections in a community teaching hospital. *Am J Infect Control*. 2010 Nov;38(9):689-93. DOI: [10.1016/j.ajic.2010.05.028](https://doi.org/10.1016/j.ajic.2010.05.028)
16. Fakhri MG, Dueweke C, Meisner S, Berriel-Cass D, Savoy-Moore R, Brach N, Rey J, DeSantis L, Saravolatz LD. Effect of nurse-led multidisciplinary rounds on reducing the unnecessary use of urinary catheterization in hospitalized patients. *Infect Control Hosp Epidemiol*. 2008 Sep;29(9):815-9. DOI: [10.1086/589584](https://doi.org/10.1086/589584)
 17. Chen YY, Chi MM, Chen YC, Chan YJ, Chou SS, Wang FD. Using a criteria-based reminder to reduce use of indwelling urinary catheters and decrease urinary tract infections. *Am J Crit Care*. 2013 Mar;22(2):105-14. DOI: [10.4037/ajcc2013464](https://doi.org/10.4037/ajcc2013464)
 18. Cornia PB, Amory JK, Fraser S, Saint S, Lipsky BA. Computer-based order entry decreases duration of indwelling urinary catheterization in hospitalized patients. *Am J Med*. 2003 Apr 1;114(5):404-7. DOI: [10.1016/S0002-9343\(02\)01568-1](https://doi.org/10.1016/S0002-9343(02)01568-1)
 19. Nagle D, Curran T, Anez-Bustillos L, Anez-Bustillo L, Poylin V. Reducing urinary tract infections in colon and rectal surgery. *Dis Colon Rectum*. 2014 Jan;57(1):91-7. DOI: [10.1097/DCR.0000000000000019](https://doi.org/10.1097/DCR.0000000000000019)
 20. Topal J, Conklin S, Camp K, Morris V, Balcezak T, Herbert P. Prevention of nosocomial catheter-associated urinary tract infections through computerized feedback to physicians and a nurse-directed protocol. *Am J Med Qual*. 2005 May-Jun;20(3):121-6. DOI: [10.1177/1062860605276074](https://doi.org/10.1177/1062860605276074)
 21. Loeb M, Hunt D, O'Halloran K, Carusone SC, Dafoe N, Walter SD. Stop orders to reduce inappropriate urinary catheterization in hospitalized patients: a randomized controlled trial. *J Gen Intern Med*. 2008 Jun;23(6):816-20. DOI: [10.1007/s11606-008-0620-2](https://doi.org/10.1007/s11606-008-0620-2)
 22. Parry MF, Grant B, Sestovic M. Successful reduction in catheter-associated urinary tract infections: focus on nurse-directed catheter removal. *Am J Infect Control*. 2013 Dec;41(12):1178-81. DOI: [10.1016/j.ajic.2013.03.296](https://doi.org/10.1016/j.ajic.2013.03.296)
 23. Mori C. A-voiding catastrophe: implementing a nurse-driven protocol. *Medsurg Nurs*. 2014 Jan-Feb;23(1):15-21, 28.
 24. Saint S, Kaufman SR, Thompson M, Rogers MA, Chenoweth CE. A reminder reduces urinary catheterization in hospitalized patients. *Jt Comm J Qual Patient Saf*. 2005 Aug;31(8):455-62. DOI: [10.1016/S1553-7250\(05\)31059-2](https://doi.org/10.1016/S1553-7250(05)31059-2)
 25. Shimoni Z, Rodrig J, Kamma N, From P. Will more restrictive indications decrease rates of urinary catheterisation? An historical comparative study. *BMJ Open*. 2012;2(2):e000473. DOI: [10.1136/bmjopen-2011-000473](https://doi.org/10.1136/bmjopen-2011-000473)
 26. Leis JA, Corpus C, Rahmani A, Catt B, Wong BM, Callery S, Vearncombe M. Medical Directive for Urinary Catheter Removal by Nurses on General Medical Wards. *JAMA Intern Med*. 2016 Jan;176(1):113-5. DOI: [10.1001/jamainternmed.2015.6319](https://doi.org/10.1001/jamainternmed.2015.6319)
 27. Dumigan DG, Kohan CA, Reed CR, Jekel JF, Fikrig MK. Utilizing national nosocomial infection surveillance system data to improve urinary tract infection rates in three intensive-care units. *Clin Perform Qual Health Care*. 1998 Oct-Dec;6(4):172-8.
 28. Kidd EA, Stewart F, Kassis NC, Hom E, Omar MI. Urethral (indwelling or intermittent) or suprapubic routes for short-term catheterisation in hospitalised adults. *Cochrane Database Syst Rev*. 2015 Dec 10;(12):CD004203. DOI: [10.1002/14651858.CD004203.pub3](https://doi.org/10.1002/14651858.CD004203.pub3)
 29. Jamison J, Maguire S, McCann J. Catheter policies for management of long term voiding problems in adults with neurogenic bladder disorders. *Cochrane Database Syst Rev*. 2013 Nov 18;(11):CD004375. DOI: [10.1002/14651858.CD004375.pub4](https://doi.org/10.1002/14651858.CD004375.pub4)
 30. Niël-Weise BS, van den Broek PJ, da Silva EM, Silva LA. Urinary catheter policies for long-term bladder drainage. *Cochrane Database Syst Rev*. 2012 Aug 15;(8):CD004201. DOI: [10.1002/14651858.CD004201.pub3](https://doi.org/10.1002/14651858.CD004201.pub3)
 31. Hunter KF, Bharmal A, Moore KN. Long-term bladder drainage: Suprapubic catheter versus other methods: a scoping review. *Neurourol Urodyn*. 2013 Sep;32(7):944-51. DOI: [10.1002/nau.22356](https://doi.org/10.1002/nau.22356)
 32. Schwartz BC, Frenette C, Lee TC, Green L, Jayaraman D. Novel low-resource intervention reduces urinary catheter use and associated urinary tract infections: role of outcome measure bias? *Am J Infect Control*. 2015 Apr;43(4):348-53. DOI: [10.1016/j.ajic.2014.12.006](https://doi.org/10.1016/j.ajic.2014.12.006)
 33. Gould CV, Umscheid CA, Agarwal RK, Kuntz G, Pegues DA; Healthcare Infection Control Practices Advisory Committee. Guideline for prevention of catheter-associated urinary tract infections 2009. *Infect Control Hosp Epidemiol*. 2010 Apr;31(4):319-26. DOI: [10.1086/651091](https://doi.org/10.1086/651091)
 34. Hooton TM, Bradley SF, Cardenas DD, Colgan R, Geerlings SE, Rice JC, Saint S, Schaeffer AJ, Tambayah PA, Tenke P, Nicolle LE; Infectious Diseases Society of America. Diagnosis,

- prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clin Infect Dis*. 2010 Mar;50(5):625-63. DOI: [10.1086/650482](https://doi.org/10.1086/650482)
35. Lo E, Nicolle LE, Coffin SE, Gould C, Maragakis LL, Meddings J, Pegues DA, Pettis AM, Saint S, Yokoe DS. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol*. 2014 May;35(5):464-79. DOI: [10.1086/675718](https://doi.org/10.1086/675718)
 36. Loveday HP, Wilson JA, Pratt RJ, Golsorkhi M, Tingle A, Bak A, Browne J, Prieto J, Wilcox M, UK Department of Health. epic3: national evidence-based guidelines for preventing healthcare-associated infections in NHS hospitals in England. *J Hosp Infect*. 2014 Jan;86 Suppl 1:S1-70. DOI: [10.1016/S0195-6701\(13\)60012-2](https://doi.org/10.1016/S0195-6701(13)60012-2)
 37. Pickard R, Bartoletti R, Bjerklund-Johansen TE. Guidelines on Urological Infections [Internet]. 2015 [cited 2016 Sep 1]. p. 31–3. Available from: <http://uroweb.org/wp-content/uploads/EAU-Extended-Guidelines-2015-Edn..pdf>
 38. Tenke P, Kovacs B, Bjerklund Johansen TE, Matsumoto T, Tambyah PA, Naber KG. European and Asian guidelines on management and prevention of catheter-associated urinary tract infections. *Int J Antimicrob Agents*. 2008 Feb;31 Suppl 1:S68-78. DOI: [10.1016/j.ijantimicag.2007.07.033](https://doi.org/10.1016/j.ijantimicag.2007.07.033)
 39. Allepuz-Palau A, Rosselló-Urgell J, Vaqué-Rafart J, Hermosilla-Pérez E, Arribas-Llorente JL, Sánchez-Payá J, Lizán-García M; EPINE Working Group. Evolution of closed urinary drainage systems use and associated factors in Spanish hospitals. *J Hosp Infect*. 2004 Aug;57(4):332-8. DOI: [10.1016/j.jhin.2004.03.026](https://doi.org/10.1016/j.jhin.2004.03.026)
 40. Tsuchida T, Makimoto K, Ohsako S, Fujino M, Kaneda M, Miyazaki T, Fujiwara F, Sugimoto T. Relationship between catheter care and catheter-associated urinary tract infection at Japanese general hospitals: a prospective observational study. *Int J Nurs Stud*. 2008 Mar;45(3):352-61. DOI: [10.1016/j.ijnurstu.2006.10.006](https://doi.org/10.1016/j.ijnurstu.2006.10.006)
 41. Prieto J, Murphy CL, Moore KN, Fader M. Intermittent catheterisation for long-term bladder management. *Cochrane Database Syst Rev*. 2014 Sep 10;(9):CD006008. DOI: [10.1002/14651858.CD006008.pub3](https://doi.org/10.1002/14651858.CD006008.pub3)
 42. Bermingham SL, Hodgkinson S, Wright S, Hayter E, Spinks J, Pellowe C. Intermittent self catheterisation with hydrophilic, gel reservoir, and non-coated catheters: a systematic review and cost effectiveness analysis. *BMJ*. 2013 Jan 8;346:e8639. DOI: [10.1136/bmj.e8639](https://doi.org/10.1136/bmj.e8639)
 43. Li L, Ye W, Ruan H, Yang B, Zhang S, Li L. Impact of hydrophilic catheters on urinary tract infections in people with spinal cord injury: systematic review and meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil*. 2013 Apr;94(4):782-7. DOI: [10.1016/j.apmr.2012.11.010](https://doi.org/10.1016/j.apmr.2012.11.010)
 44. Lam TB, Omar MI, Fisher E, Gillies K, MacLennan S. Types of indwelling urethral catheters for short-term catheterisation in hospitalised adults. *Cochrane Database Syst Rev*. 2014 Sep 23;(9):CD004013. DOI: [10.1002/14651858.CD004013.pub4](https://doi.org/10.1002/14651858.CD004013.pub4)
 45. Pickard R, Lam T, MacLennan G, Starr K, Kilonzo M, McPherson G, Gillies K, McDonald A, Walton K, Buckley B, Glazener C, Boachie C, Burr J, Norrie J, Vale L, Grant A, N'dow J. Types of urethral catheter for reducing symptomatic urinary tract infections in hospitalised adults requiring short-term catheterisation: multicentre randomised controlled trial and economic evaluation of antimicrobial- and antiseptic-impregnated urethral catheters (the CATHETER trial). *Health Technol Assess*. 2012 Nov;16(47):1-197. DOI: [10.3310/hta16470](https://doi.org/10.3310/hta16470)
 46. Kilonzo M, Vale L, Pickard R, Lam T, N'Dow J; Catheter Trial Group. Cost effectiveness of antimicrobial catheters for adults requiring short-term catheterisation in hospital. *Eur Urol*. 2014 Oct;66(4):615-8. DOI: [10.1016/j.eururo.2014.05.035](https://doi.org/10.1016/j.eururo.2014.05.035)
 47. Jahn P, Beutner K, Langer G. Types of indwelling urinary catheters for long-term bladder drainage in adults. *Cochrane Database Syst Rev*. 2012 Oct 17;10:CD004997. DOI: [10.1002/14651858.CD004997.pub3](https://doi.org/10.1002/14651858.CD004997.pub3)
 48. Lusardi G, Lipp A, Shaw C. Antibiotic prophylaxis for short-term catheter bladder drainage in adults. *Cochrane Database Syst Rev*. 2013 Jul 3;(7):CD005428. DOI: [10.1002/14651858.CD005428.pub2](https://doi.org/10.1002/14651858.CD005428.pub2)
 49. Morton SC, Shekelle PG, Adams JL, Bennett C, Dobkin BH, Montgomerie J, Vickrey BG. Antimicrobial prophylaxis for urinary tract infection in persons with spinal cord dysfunction. *Arch Phys Med Rehabil*. 2002 Jan;83(1):129-38.
 50. Marschall J, Carpenter CR, Fowler S, Trautner BW; CDC Prevention Epicenters Program.

- Antibiotic prophylaxis for urinary tract infections after removal of urinary catheter: meta-analysis. *BMJ*. 2013 Jun 11;346:f3147. DOI: [10.1136/bmj.f3147](https://doi.org/10.1136/bmj.f3147)
51. Maki DG. ACP Journal Club. Review: Antibiotic prophylaxis on removal of urinary catheters reduces symptomatic urinary tract infections. *Ann Intern Med*. 2013 Oct 15;159(8):JC9. DOI: [10.7326/0003-4819-159-8-201310150-02009](https://doi.org/10.7326/0003-4819-159-8-201310150-02009)
 52. MacFadden DR, Ridgway JP, Robicsek A, Elligsen M, Daneman N. Predictive utility of prior positive urine cultures. *Clin Infect Dis*. 2014 Nov;59(9):1265-71. DOI: [10.1093/cid/ciu588](https://doi.org/10.1093/cid/ciu588)
 53. Darouiche RO, Al Mohajer M, Siddiq DM, Minard CG. Short versus long course of antibiotics for catheter-associated urinary tract infections in patients with spinal cord injury: a randomized controlled noninferiority trial. *Arch Phys Med Rehabil*. 2014 Feb;95(2):290-6. DOI: [10.1016/j.apmr.2013.09.003](https://doi.org/10.1016/j.apmr.2013.09.003)
 54. Warren JW, Anthony WC, Hoopes JM, Muncie HL Jr. Cephalexin for susceptible bacteriuria in afebrile, long-term catheterized patients. *JAMA*. 1982 Jul;248(4):454-8.
 55. Klevens RM, Edwards JR, Richards CL Jr, Horan TC, Gaynes RP, Pollock DA, Cardo DM. Estimating health care-associated infections and deaths in U.S. hospitals, 2002. *Public Health Rep*. 2007 Mar-Apr;122(2):160-6. DOI: [10.1177/003335490712200205](https://doi.org/10.1177/003335490712200205)
 56. Weinstein JW, Mazon D, Pantelick E, Reagan-Cirincione P, Dembry LM, Hierholzer WJ Jr. A decade of prevalence surveys in a tertiary-care center: trends in nosocomial infection rates, device utilization, and patient acuity. *Infect Control Hosp Epidemiol*. 1999 Aug;20(8):543-8. DOI: [10.1086/501675](https://doi.org/10.1086/501675)
 57. Centers for Disease Control and Prevention. Catheter-associated urinary tract infections [Internet]. [cited 2016 Sep 2]. Available from: http://www.cdc.gov/HAI/ca_uti/uti.html
 58. Gould CV, Umscheid CA, Agarwal RK, Kuntz G, Pegues DA; Healthcare Infection Control Practices Advisory Committee. Guideline for prevention of catheter-associated urinary tract infections 2009. *Infect Control Hosp Epidemiol*. 2010 Apr;31(4):319-26. DOI: [10.1086/651091](https://doi.org/10.1086/651091)
 59. Umscheid CA, Mitchell MD, Doshi JA, Agarwal R, Williams K, Brennan PJ. Estimating the proportion of healthcare-associated infections that are reasonably preventable and the related mortality and costs. *Infect Control Hosp Epidemiol*. 2011 Feb;32(2):101-14. DOI: [10.1086/657912](https://doi.org/10.1086/657912)
 60. Yi SH, Baggs J, Gould CV, Scott RD 2nd, Jernigan JA. Medicare reimbursement attributable to catheter-associated urinary tract infection in the inpatient setting: a retrospective cohort analysis. *Med Care*. 2014 Jun;52(6):469-78. DOI: [10.1097/MLR.000000000000106](https://doi.org/10.1097/MLR.000000000000106)
 61. Hu B, Tao L, Rosenthal VD, Liu K, Yun Y, Suo Y, Gao X, Li R, Su D, Wang H, Hao C, Pan W, Saunders CL. Device-associated infection rates, device use, length of stay, and mortality in intensive care units of 4 Chinese hospitals: International Nosocomial Control Consortium findings. *Am J Infect Control*. 2013 Apr;41(4):301-6. DOI: [10.1016/j.ajic.2012.03.037](https://doi.org/10.1016/j.ajic.2012.03.037)
 62. Castle N, Ferguson-Rome JC, Teresi JA. Elder abuse in residential long-term care: an update to the 2003 National Research Council report. *J Appl Gerontol*. 2015 Jun;34(4):407-43. DOI: [10.1177/0733464813492583](https://doi.org/10.1177/0733464813492583)
 63. Ling ML, Apisarnthanarak A, Madriaga G. The Burden of Healthcare-Associated Infections in Southeast Asia: A Systematic Literature Review and Meta-analysis. *Clin Infect Dis*. 2015 Jun;60(11):1690-9. DOI: [10.1093/cid/civ095](https://doi.org/10.1093/cid/civ095)
 64. Maki DG, Tambyah PA. Engineering out the risk for infection with urinary catheters. *Emerg Infect Dis*. 2001 Mar-Apr;7(2):342-7.
 65. Nicolle LE. Urinary catheter-associated infections. *Infect Dis Clin North Am*. 2012 Mar;26(1):13-27. DOI: [10.1016/j.idc.2011.09.009](https://doi.org/10.1016/j.idc.2011.09.009)
 66. Foxman B. The epidemiology of urinary tract infection. *Nat Rev Urol*. 2010 Dec;7(12):653-60. DOI: [10.1038/nrurol.2010.190](https://doi.org/10.1038/nrurol.2010.190)
 67. Platt R, Polk BF, Murdock B, Rosner B. Risk factors for nosocomial urinary tract infection. *Am J Epidemiol*. 1986 Dec;124(6):977-85. DOI: [10.1093/oxfordjournals.aje.a114487](https://doi.org/10.1093/oxfordjournals.aje.a114487)
 68. Barbadoro P, Labricciosa FM, Recanatini C, Gori G, Tirabassi F, Martini E, Gioia MG, D'Errico MM, Prospero E. Catheter-associated urinary tract infection: Role of the setting of catheter insertion. *Am J Infect Control*. 2015 Jul;43(7):707-10. DOI: [10.1016/j.ajic.2015.02.011](https://doi.org/10.1016/j.ajic.2015.02.011)
 69. Lewis SS, Knelson LP, Moehring RW, Chen LF, Sexton DJ, Anderson DJ. Comparison of non-intensive care unit (ICU) versus ICU rates of catheter-associated urinary tract infection in community hospitals. *Infect Control Hosp Epidemiol*. 2013 Jul;34(7):744-7. DOI: [10.1086/671000](https://doi.org/10.1086/671000)
 70. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, Bernstein SJ. The Ann Arbor Criteria for Appropriate Urinary Catheter Use in Hospitalized Medical Patients: Results Obtained

- by Using the RAND/UCLA Appropriateness Method. *Ann Intern Med.* 2015 May;162(9 Suppl):S1-34. DOI: [10.7326/M14-1304](https://doi.org/10.7326/M14-1304)
71. Griffiths R, Fernandez R. Strategies for the removal of short-term indwelling urethral catheters in adults. *Cochrane Database Syst Rev.* 2007 Apr 18;(2):CD004011. DOI: [10.1002/14651858.CD004011.pub3](https://doi.org/10.1002/14651858.CD004011.pub3)
 72. Dixon L, Dolan LM, Brown K, Hilton P. RCT of urethral versus suprapubic catheterization. *Br J Nurs.* 2010 Oct 14-27;19(18):S7-13. DOI: [10.12968/bjon.2010.19.Sup8.79063](https://doi.org/10.12968/bjon.2010.19.Sup8.79063)
 73. Patel MI, Watts W, Grant A. The optimal form of urinary drainage after acute retention of urine. *BJU Int.* 2001 Jul;88(1):26-9.
 74. Beuscher T. Pad weighing for reduction of indwelling urinary use and catheter-associated urinary tract infection: a quality improvement project. *J Wound Ostomy Continence Nurs.* 2014 Nov-Dec;41(6):604-8. DOI: [10.1097/WON.0000000000000068](https://doi.org/10.1097/WON.0000000000000068)
 75. Huth TS, Burke JP, Larsen RA, Classen DC, Stevens LE. Clinical trial of junction seals for the prevention of urinary catheter-associated bacteriuria. *Arch Intern Med.* 1992 Apr;152(4):807-12. DOI: [10.1001/archinte.1992.00400160103019](https://doi.org/10.1001/archinte.1992.00400160103019)
 76. DeGroot-Kosolcharoen J, Guse R, Jones JM. Evaluation of a urinary catheter with a preconnected closed drainage bag. *Infect Control Hosp Epidemiol.* 1988 Feb;9(2):72-6. DOI: [10.2307/30144145](https://doi.org/10.2307/30144145)
 77. Classen DC, Larsen RA, Burke JP, Stevens LE. Prevention of catheter-associated bacteriuria: clinical trial of methods to block three known pathways of infection. *Am J Infect Control.* 1991 Jun;19(3):136-42. DOI: [10.1016/0196-6553\(91\)90019-9](https://doi.org/10.1016/0196-6553(91)90019-9)
 78. Webster J, Hood RH, Burrige CA, Doidge ML, Phillips KM, George N. Water or antiseptic for periurethral cleaning before urinary catheterization: a randomized controlled trial. *Am J Infect Control.* 2001 Dec;29(6):389-94. DOI: [10.1067/mic.2001.117447](https://doi.org/10.1067/mic.2001.117447)
 79. Cheung K, Leung P, Wong YC, To OK, Yeung YF, Chan MW, Yip YL, Kwok CW. Water versus antiseptic periurethral cleansing before catheterization among home care patients: a randomized controlled trial. *Am J Infect Control.* 2008 Jun;36(5):375-80. DOI: [10.1016/j.ajic.2007.03.004](https://doi.org/10.1016/j.ajic.2007.03.004)
 80. Carapeti EA, Andrews SM, Bentley PG. Randomised study of sterile versus non-sterile urethral catheterisation. *Ann R Coll Surg Engl.* 1996 Jan;78(1):59-60.
 81. Darouiche RO, Goetz L, Kaldis T, Cerra-Stewart C, AlSharif A, Priebe M. Impact of StatLock securing device on symptomatic catheter-related urinary tract infection: a prospective, randomized, multicenter clinical trial. *Am J Infect Control.* 2006 Nov;34(9):555-60. DOI: [10.1016/j.ajic.2006.03.010](https://doi.org/10.1016/j.ajic.2006.03.010)
 82. Hagen S, Sinclair L, Cross S. Washout policies in long-term indwelling urinary catheterisation in adults. *Cochrane Database Syst Rev.* 2010 Mar 17;(3):CD004012. DOI: [10.1002/14651858.CD004012.pub4](https://doi.org/10.1002/14651858.CD004012.pub4)
 83. Waites KB, Canupp KC, Roper JF, Camp SM, Chen Y. Evaluation of 3 methods of bladder irrigation to treat bacteriuria in persons with neurogenic bladder. *J Spinal Cord Med.* 2006;29(3):217-26. DOI: [10.1080/10790268.2006.11753877](https://doi.org/10.1080/10790268.2006.11753877)
 84. Willson M, Wilde M, Webb ML, Thompson D, Parker D, Harwood J, Callan L, Gray M. Nursing interventions to reduce the risk of catheter-associated urinary tract infection: part 2: staff education, monitoring, and care techniques. *J Wound Ostomy Continence Nurs.* 2009 Mar-Apr;36(2):137-54. DOI: [10.1097/01.WON.0000347655.56851.04](https://doi.org/10.1097/01.WON.0000347655.56851.04)
 85. Burke JP, Garibaldi RA, Britt MR, Jacobson JA, Conti M, Alling DW. Prevention of catheter-associated urinary tract infections. Efficacy of daily meatal care regimens. *Am J Med.* 1981 Mar;70(3):655-8. DOI: [10.1016/0002-9343\(81\)90591-X](https://doi.org/10.1016/0002-9343(81)90591-X)
 86. Huth TS, Burke JP, Larsen RA, Classen DC, Stevens LE. Randomized trial of meatal care with silver sulfadiazine cream for the prevention of catheter-associated bacteriuria. *J Infect Dis.* 1992 Jan;165(1):14-8. DOI: [10.1093/infdis/165.1.14](https://doi.org/10.1093/infdis/165.1.14)
 87. Strouse AC. Appraising the Literature On Bathing Practices And Catheter-Associated Urinary Tract Infection Prevention. *Urol Nurs.* 2015 Jan-Feb;35(1):11-7.
 88. Evans HL, Dellit TH, Chan J, Nathens AB, Maier RV, Cuschieri J. Effect of chlorhexidine whole-body bathing on hospital-acquired infections among trauma patients. *Arch Surg.* 2010 Mar;145(3):240-6. DOI: [10.1001/archsurg.2010.5](https://doi.org/10.1001/archsurg.2010.5)
 89. Bleasdale SC, Trick WE, Gonzalez IM, Lyles RD, Hayden MK, Weinstein RA. Effectiveness of chlorhexidine bathing to reduce catheter-associated bloodstream infections in medical intensive care unit patients. *Arch Intern Med.* 2007 Oct;167(19):2073-9. DOI: [10.1001/archinte.167.19.2073](https://doi.org/10.1001/archinte.167.19.2073)

90. Leone M, Garnier F, Antonini F, Bimar MC, Albanèse J, Martin C. Comparison of effectiveness of two urinary drainage systems in intensive care unit: a prospective, randomized clinical trial. *Intensive Care Med.* 2003 Mar;29(3):410-3. DOI: [10.1007/s00134-003-1644-z](https://doi.org/10.1007/s00134-003-1644-z)
91. Panitchote A, Charoensri S, Chetchotisakd P, Hurst C. Pilot study of a non-return catheter valve for reducing catheter-associated urinary tract infections in critically ill patients. *J Med Assoc Thai.* 2015 Feb;98(2):150-5.
92. Cooper FP, Alexander CE, Sinha S, Omar MI. Policies for replacing long-term indwelling urinary catheters in adults. *Cochrane Database Syst Rev.* 2016 Jul 26;7:CD011115. DOI: [10.1002/14651858.CD011115.pub2](https://doi.org/10.1002/14651858.CD011115.pub2)
93. Hachem R, Reitzel R, Borne A, Jiang Y, Tinkey P, Uthamanthil R, Chandra J, Ghannoum M, Raad I. Novel antiseptic urinary catheters for prevention of urinary tract infections: correlation of in vivo and in vitro test results. *Antimicrob Agents Chemother.* 2009 Dec;53(12):5145-9. DOI: [10.1128/AAC.00718-09](https://doi.org/10.1128/AAC.00718-09)
94. Shapur NK, Duvdevani M, Friedman M, Zaks B, Gati I, Lavy E, Katz R, Landau EH, Pode D, Gofrit ON, Steinberg D. Sustained release varnish containing chlorhexidine for prevention of biofilm formation on urinary catheter surface: in vitro study. *J Endourol.* 2012 Jan;26(1):26-31. DOI: [10.1089/end.2011.0140](https://doi.org/10.1089/end.2011.0140)
95. Segev G, Bankirer T, Steinberg D, Duvdevani M, Shapur NK, Friedman M, Lavy E. Evaluation of urinary catheters coated with sustained-release varnish of chlorhexidine in mitigating biofilm formation on urinary catheters in dogs. *J Vet Intern Med.* 2013 Jan-Feb;27(1):39-46. DOI: [10.1111/j.1939-1676.2012.01027.x](https://doi.org/10.1111/j.1939-1676.2012.01027.x)
96. Amalaradjou MA, Narayanan A, Baskaran SA, Venkitanarayanan K. Antibiofilm effect of trans-cinnamaldehyde on uropathogenic *Escherichia coli*. *J Urol.* 2010 Jul;184(1):358-63. DOI: [10.1016/j.juro.2010.03.006](https://doi.org/10.1016/j.juro.2010.03.006)
97. Evliyaoğlu Y, Kobaner M, Celebi H, Yelsel K, Doğan A. The efficacy of a novel antibacterial hydroxyapatite nanoparticle-coated indwelling urinary catheter in preventing biofilm formation and catheter-associated urinary tract infection in rabbits. *Urol Res.* 2011 Dec;39(6):443-9. DOI: [10.1007/s00240-011-0379-5](https://doi.org/10.1007/s00240-011-0379-5)
98. Williams GJ, Stickler DJ. Effect of triclosan on the formation of crystalline biofilms by mixed communities of urinary tract pathogens on urinary catheters. *J Med Microbiol.* 2008 Sep;57(Pt 9):1135-40. DOI: [10.1099/jmm.0.2008/002295-0](https://doi.org/10.1099/jmm.0.2008/002295-0)
99. Sun Y, Zeng Q, Zhang Z, Xu C, Wang Y, He J. Decreased urethral mucosal damage and delayed bacterial colonization during short-term urethral catheterization using a novel trefoil urethral catheter profile in rabbits. *J Urol.* 2011 Oct;186(4):1497-501. DOI: [10.1016/j.juro.2011.05.043](https://doi.org/10.1016/j.juro.2011.05.043)
100. Syed MA, Manzoor U, Shah I, Bukhari SH. Antibacterial effects of Tungsten nanoparticles on the *Escherichia coli* strains isolated from catheterized urinary tract infection (UTI) cases and *Staphylococcus aureus*. *New Microbiol.* 2010 Oct;33(4):329-35.
101. Liao KS, Lehman SM, Twardy DJ, Donlan RM, Trautner BW. Bacteriophages are synergistic with bacterial interference for the prevention of *Pseudomonas aeruginosa* biofilm formation on urinary catheters. *J Appl Microbiol.* 2012 Dec;113(6):1530-9. DOI: [10.1111/j.1365-2672.2012.05432.x](https://doi.org/10.1111/j.1365-2672.2012.05432.x)
102. Prasad A, Cevallos ME, Riosa S, Darouiche RO, Trautner BW. A bacterial interference strategy for prevention of UTI in persons practicing intermittent catheterization. *Spinal Cord.* 2009 Jul;47(7):565-9. DOI: [10.1038/sc.2008.166](https://doi.org/10.1038/sc.2008.166)
103. Trautner BW, Hull RA, Thornby JI, Darouiche RO. Coating urinary catheters with an avirulent strain of *Escherichia coli* as a means to establish asymptomatic colonization. *Infect Control Hosp Epidemiol.* 2007 Jan;28(1):92-4. DOI: [10.1086/510872](https://doi.org/10.1086/510872)
104. Tenke P, Köves B, Nagy K, Hultgren SJ, Mendling W, Wullt B, Grabe M, Wagenlehner FM, Cek M, Pickard R, Botto H, Naber KG, Bjerklund Johansen TE. Update on biofilm infections in the urinary tract. *World J Urol.* 2012 Feb;30(1):51-7. DOI: [10.1007/s00345-011-0689-9](https://doi.org/10.1007/s00345-011-0689-9)
105. Shenderovich J, Feldman M, Kirmayer D, Al-Quntar A, Steinberg D, Lavy E, Friedman M. Local sustained-release delivery systems of the antibiofilm agent thiazolidinedione-8 for prevention of catheter-associated urinary tract infections. *Int J Pharm.* 2015 May;485(1-2):164-70. DOI: [10.1016/j.ijpharm.2015.02.067](https://doi.org/10.1016/j.ijpharm.2015.02.067)
106. Ivanova K, Fernandes MM, Mendoza E, Tzanov T. Enzyme multilayer coatings inhibit *Pseudomonas aeruginosa* biofilm formation on urinary catheters. *Appl Microbiol Biotechnol.* 2015 May;99(10):4373-85. DOI: [10.1007/s00253-015-6378-7](https://doi.org/10.1007/s00253-015-6378-7)

107. Schumm K, Lam TB. Types of urethral catheters for management of short-term voiding problems in hospitalised adults. *Cochrane Database Syst Rev.* 2008 Apr 16;(2):CD004013. DOI: [10.1002/14651858.CD004013.pub3](https://doi.org/10.1002/14651858.CD004013.pub3)
108. Pickard R, Lam T, MacLennan G, Starr K, Kilonzo M, McPherson G, Gillies K, McDonald A, Walton K, Buckley B, Glazener C, Boachie C, Burr J, Norrie J, Vale L, Grant A, N'Dow J. Antimicrobial catheters for reduction of symptomatic urinary tract infection in adults requiring short-term catheterisation in hospital: a multicentre randomised controlled trial. *Lancet.* 2012 Dec 1;380(9857):1927-35. DOI: [10.1016/S0140-6736\(12\)61380-4](https://doi.org/10.1016/S0140-6736(12)61380-4)
109. Makuta G, Chrysaifis M, Lam T. Measuring the efficacy of antimicrobial catheters. *Nurs Times.* 2013 Nov 6-12;109(44):16, 18-9.
110. Wolters HH, Palmes D, Lordugin E, Bahde R, Senninger N, Hölzen JP, Kobschull L. Antibiotic prophylaxis at urinary catheter removal prevents urinary tract infection after kidney transplantation. *Transplant Proc.* 2014 Dec;46(10):3463-5. DOI: [10.1016/j.transproceed.2014.04.019](https://doi.org/10.1016/j.transproceed.2014.04.019)
111. Salomon J, Denys P, Merle C, Chartier-Kastler E, Perronne C, Gaillard JL, Bernard L. Prevention of urinary tract infection in spinal cord-injured patients: safety and efficacy of a weekly oral cyclic antibiotic (WOCA) programme with a 2 year follow-up--an observational prospective study. *J Antimicrob Chemother.* 2006 Apr;57(4):784-8. DOI: [10.1093/jac/dkl010](https://doi.org/10.1093/jac/dkl010)
112. McMurdo ME, Bissett LY, Price RJ, Phillips G, Crombie IK. Does ingestion of cranberry juice reduce symptomatic urinary tract infections in older people in hospital? A double-blind, placebo-controlled trial. *Age Ageing.* 2005 May;34(3):256-61. DOI: [10.1093/ageing/afi101](https://doi.org/10.1093/ageing/afi101)
113. Opperman EA. Cranberry is not effective for the prevention or treatment of urinary tract infections in individuals with spinal cord injury. *Spinal Cord.* 2010 Jun;48(6):451-6. DOI: [10.1038/sc.2009.159](https://doi.org/10.1038/sc.2009.159)
114. Chung YC, Chen HH, Yeh ML. Vinegar for decreasing catheter-associated bacteriuria in long-term catheterized patients: a randomized controlled trial. *Biol Res Nurs.* 2012 Jul;14(3):294-301. DOI: [10.1177/1099800411412767](https://doi.org/10.1177/1099800411412767)
115. Lee BS, Bhuta T, Simpson JM, Craig JC. Methenamine hippurate for preventing urinary tract infections. *Cochrane Database Syst Rev.* 2012 Oct 17;10:CD003265. DOI: [10.1002/14651858.CD003265.pub3](https://doi.org/10.1002/14651858.CD003265.pub3)
116. Darouiche RO, Green BG, Donovan WH, Chen D, Schwartz M, Merritt J, Mendez M, Hull RA. Multicenter randomized controlled trial of bacterial interference for prevention of urinary tract infection in patients with neurogenic bladder. *Urology.* 2011 Aug;78(2):341-6. DOI: [10.1016/j.urology.2011.03.062](https://doi.org/10.1016/j.urology.2011.03.062)
117. Hull R, Rudy D, Donovan W, Svanborg C, Wieser I, Stewart C, Darouiche R. Urinary tract infection prophylaxis using *Escherichia coli* 83972 in spinal cord injured patients. *J Urol.* 2000 Mar;163(3):872-7. DOI: [10.1016/S0022-5347\(05\)67823-8](https://doi.org/10.1016/S0022-5347(05)67823-8)
118. Harding GK, Nicolle LE, Ronald AR, Preiksaitis JK, Forward KR, Low DE, Cheang M. How long should catheter-acquired urinary tract infection in women be treated? A randomized controlled study. *Ann Intern Med.* 1991 May;114(9):713-9. DOI: [10.7326/0003-4819-114-9-713](https://doi.org/10.7326/0003-4819-114-9-713)
119. Dow G, Rao P, Harding G, Brunka J, Kennedy J, Alfa M, Nicolle LE. A prospective, randomized trial of 3 or 14 days of ciprofloxacin treatment for acute urinary tract infection in patients with spinal cord injury. *Clin Infect Dis.* 2004 Sep;39(5):658-64. DOI: [10.1086/423000](https://doi.org/10.1086/423000)
120. Leone M, Perrin AS, Granier I, Visintini P, Blasco V, Antonini F, Albanèse J, Martin C. A randomized trial of catheter change and short course of antibiotics for asymptomatic bacteriuria in catheterized ICU patients. *Intensive Care Med.* 2007 Apr;33(4):726-9. DOI: [10.1007/s00134-007-0534-1](https://doi.org/10.1007/s00134-007-0534-1)

Corresponding authors: Professor Paul Anantharajah Tambyah, Yong Loo Lin School of Medicine, National University Hospital, Department of Medicine, 1E Kent Ridge Road, 119228 Singapore, Singapore, Phone: -, E-mail: paul_anantharajah_tambyah@nuhs.edu.sg

Citation note: Tambyah PA, Chuang L, Chiong E. Prevention of catheter-associated UTI. Version: 2018-08-06. In: Bjerklund Johansen TE, Wagenlehner FME, Matsumoto T, Cho YH, Krieger JN, Shoskes D, Naber KG, editors. *Urogenital Infections and Inflammations*. Berlin: German Medical Science GMS Publishing House; 2017-. DOI: [10.5680/lhuiu000030](https://doi.org/10.5680/lhuiu000030)

Copyright: © 2025 Paul Anantharajah Tambyah et al.
This is an Open Access publication distributed under the terms of the Creative Commons Attribution 4.0 International License. See license information at <https://creativecommons.org/licenses/by/4.0/>